



University of North Dakota
UND Scholarly Commons

Education, Health & Behavior Studies Faculty
Publications

Department of Education, Health & Behavior
Studies

2-2016

Australia and Other Nations Are Failing to Meet Sedentary Behaviour Guidelines for Children: Implications and a Way Forward

Leon Straker

Erin Kaye Howie

Dylan Paul Cliff

Melanie T. Davern

Lina Engelen

See next page for additional authors

Follow this and additional works at: <https://commons.und.edu/ehb-fac>



Part of the [Health and Physical Education Commons](#)

Recommended Citation

Straker, Leon; Howie, Erin Kaye; Cliff, Dylan Paul; Davern, Melanie T.; Engelen, Lina; Gomersall, Sjaan R.; Ziviani, Jenny; Schranz, Natasha K.; Olds, Tim; and Tomkinson, Grant, "Australia and Other Nations Are Failing to Meet Sedentary Behaviour Guidelines for Children: Implications and a Way Forward" (2016). *Education, Health & Behavior Studies Faculty Publications*. 15.

<https://commons.und.edu/ehb-fac/15>

This Article is brought to you for free and open access by the Department of Education, Health & Behavior Studies at UND Scholarly Commons. It has been accepted for inclusion in Education, Health & Behavior Studies Faculty Publications by an authorized administrator of UND Scholarly Commons. For more information, please contact zeineb.yousif@library.und.edu.

Authors

Leon Straker, Erin Kaye Howie, Dylan Paul Cliff, Melanie T. Davern, Lina Engelen, Sjaan R. Gomersall, Jenny Ziviani, Natasha K. Schranz, Tim Olds, and Grant Tomkinson

Australia and Other Nations Are Failing to Meet Sedentary Behaviour Guidelines for Children: Implications and a Way Forward

Leon Straker, Erin Kaye Howie, Dylan Paul Cliff, Melanie T. Davern, Lina Engelen, Sjaan R. Gomersall, Jenny Ziviani, Natasha K. Schranz, Tim Olds, and Grant Ryan Tomkinson

Background: Australia has joined a growing number of nations that have evaluated the physical activity and sedentary behavior status of their children. Australia received a “D minus” in the first Active Healthy Kids Australia Physical Activity Report Card. **Methods:** An expert subgroup of the Australian Report Card Research Working Group iteratively reviewed available evidence to answer 3 questions: (a) What are the main sedentary behaviors of children? (b) What are the potential mechanisms for sedentary behavior to impact child health and development? and (c) What are the effects of different types of sedentary behaviors on child health and development? **Results:** Neither sedentary time nor screen time is a homogeneous activity likely to result in homogenous effects. There are several mechanisms by which various sedentary behaviors may positively or negatively affect cardiometabolic, neuromusculoskeletal, and psychosocial health, though the strength of evidence varies. National surveillance systems and mechanistic, longitudinal, and experimental studies are needed for Australia and other nations to improve their grade. **Conclusions:** Despite limitations, available evidence is sufficiently convincing that the total exposure and pattern of exposure to sedentary behaviors are critical to the healthy growth, development, and wellbeing of children. Nations therefore need strategies to address these common behaviors.

Keywords: screen time, child health, well-being, sedentary behaviour, indicators

In May 2014, 15 countries gathered in Toronto, Canada, for the Global Summit on Physical Activity of Children in response to international concern over the physical inactivity of the world’s children. Using expert consensus panels, countries reviewed their respective available data and weighed the evidence to assign a grade for 9 core indicators in national Physical Activity Report Cards. The core indicators were related to individual behaviors that contributed to overall physical activity levels, as well as sources of influence and strategies and investments. One of the core behavioral indicators was sedentary behavior, which was operationalized as the proportion of children and young people meeting the recommended national screen time guidelines. For Australia, this is spending no more than 1 hour per day for 2- to 4-year-olds and less than 2 hours per day for 5- to 17-year-olds viewing an electronic screen for leisure purposes.¹ Currently there are no national data for children less than 2 years of age to determine what percentage are complying with the national guideline of no screen time.

Australia received a grade of “D minus(–)” for sedentary behaviors, with only 29% of 5- to 17-year-olds meeting screen time recommendations.^{2,3} Fewer Australian teenagers met the recommendations (19% of 15- to 17-year-olds) than younger school children (41% of 5- to 8-year-olds and 24% of 9- to 14-year-olds) or preschoolers (26% of 2- to 4-year-olds).³ Australia is not alone, with 4 other countries rated below Australia with a “Fail” and 4 more with a “D” in sedentary behavior. The highest grade achieved was a “B,” by Ghana and Kenya, followed by New Zealand and Ireland which both received grades of “C” (see Table 1). While the metrics used to assign grades varied between countries, the grades assigned raise the question: What can countries do to improve their grades?

Australia’s sedentary behavior grade was based on the percentage of children meeting the recommendations for daily screen time, as it generally was for other nations (though the exact definitions varied). The Active Healthy Kids Australia Physical Activity Report Card focused on screen time sedentary behavior for a number of reasons. First, national guidelines recommend a dose specifically for screen-based sedentary behaviors,¹ and the best nationally representative data available in Australia were

for compliance with screen time guidelines rather than all sedentary behaviors. Secondly, the Research Working Group (24 experts in the field of physical activity and health from around Australia who evaluated the evidence and assigned a grade by consensus) had more confidence in reported screen time than other self- or proxy- report measures of sedentary behaviors.⁴ Thirdly, there was stronger evidence that screen time, particularly television (TV) watching, was associated with detrimental outcomes (see Question 3 section for further details⁵). However, basing the grade solely on meeting screen time guidelines is a limitation for multiple reasons: (a) much of childhood sedentary behavior is not screen-based; (b) overall sedentary behavior, in addition to screen time, potentially has detrimental effects^{6,7}; and (c) screen time itself is varied and changing rapidly.

Methods

The following is a discussion of key evidence that resulted from a critical review by an expert subgroup of the Australian Report Card Research Working Group. The Research Working Group had been

Table 1 Summary of Sedentary Behavior Grades in National Physical Activity Report Cards

| Country | Grade | Percentage (%) meeting screen time guidelines (≤ 2 hours/day unless otherwise noted)* |
|---------------|-------|------------------------------------------------------------------------------------------------|
| Ghana | B | 79% of 13–17-year-olds (global physical activity guidelines, < 3 hours sitting) |
| Kenya | B | Average of 1.75 hours screen time on school day, 4.25 hours on weekend days for 9–11-year-olds |
| New Zealand | C | 60% of 5–9-year-olds, 33% of 10–14-year-olds |
| Ireland | C– | 54% of 11–15-year-olds (TV only) |
| Colombia | D | 42% of 5–12-year-olds |
| Finland | D | 22% of 11–15-year-olds (on weekdays) |
| Mexico | D | 33% of 10–18-year-olds |
| United States | D | 59% of 6–8-year-olds, 48% of 9–11-year-olds (but ethnic disparities) |
| Australia | D– | 29% of 5–17-year-olds |
| Canada | F | 69% of 5–11-year-olds, 19% of 10–16-year-olds |
| Nigeria | F | 5–35% of 6–18-year-olds (< 3 hours per day) |
| Scotland | F | 24% of 11–15-year-olds (TV only) |
| South Africa | F | Average 3 hours TV per day for 10–17-year-olds |

Note. Estimates are taken from respective country report cards, and the definitions of meeting guidelines varied, as did the survey instruments used and age groups assessed.

collecting and evaluating literature and data related to the Report Card generation. To conduct the present review, the first 2 authors conducted a further literature search of primary databases to capture recently published evidence. The critical analysis followed an iterative process by the expert subgroup where additional literature was considered and all evidence was synthesized. The experts reviewed the literature in reference to 3 general questions about sedentary behaviors, as seen in Figure 1. A better understanding of the answers to these 3 questions will help inform strategies to reduce sedentary behaviors among children and thus improve the grade.

Question 1: What Are the Main Sedentary Behaviors of Children?

Sedentary behavior is defined as any waking behavior with a low energy expenditure (< 1.5 metabolic equivalents [METs]) and a sitting or reclined posture⁸ and is part of a spectrum of “activity” of various energy expenditure intensities ranging from sedentary, through light (typically ≥ 1.5 – < 3 METs) to moderate (≥ 3 – < 6 METs) and vigorous (≥ 6 METs). Although there has been debate on the specific MET cutpoints used for children,⁹ research in young children suggests that 1.5 METs is consistent with the energy cost of sedentary activities.¹⁰ Thus each child’s 24-hour day can be divided into sleep and wake “activity,” with “activity” further divided by intensity into sedentary, light, moderate, and vigorous time. The most common measures of sedentary behavior are self-report and accelerometry, which both have limitations.¹¹ Self- or proxy-report questionnaires and recalls are subject to recall bias, and some continue to show limited validity compared with device-based or objective measures, and accelerometers do not distinguish between types of sedentary behaviors or provide context. Inclinometers have been increasingly used to measure sedentary time as they better distinguish between postures of sedentary

behaviors (ie, lying, sitting, standing), but still do not provide context or type of behavior. Accelerometers can yield widely discrepant estimates of sedentary time according to device placement and analytical decisions around nonwear time, operationalization of sleep, epoch length, and intensity cut-offs. This is only a brief description of some of the issues surrounding the measurement of sedentary behaviors in children, a topic that warrants further discussion beyond this review.

Being sedentary is seen as different to not attaining recommended daily amounts of moderate to vigorous physical activity (MVPA), as a child can spend a large portion of his or her day in sedentary behavior but still meet daily MVPA recommendations of at least 60 minutes.¹² Further, the health effects of accumulating too little physical activity or too much sedentary time may differ,^{13–15} although the research evidence in children is still building.^{16–19}

The largest proportion of a child's waking day is spent in sedentary behavior. For example, accelerometer data on Australian 10- to 12-year-olds showed that 63% of their waking day was spent engaged in sedentary activities, as shown in Figure 2.20. While objective surveillance of Australian children's physical activity is limited, studies suggest that preschool-aged children,^{21,22} primary school aged children²³ and young adolescents²⁴ spend at least 60% of wake time in sedentary behaviors, which is consistent with data from 39 countries.²⁵ These data also suggest that the proportion of the waking day spent sedentary increases with age across childhood, although the evidence for young children and how sedentary behavior tracks throughout childhood into adulthood is limited.²⁶

Sedentary behavior can be thought to occur in 4 main domains of children's lives—education/school/child care, transport, self-care/ domestic chores, and leisure/play. For school-aged children, a main “occupation” is that of being a student in which the majority of time at school is sedentary.²⁰ Educational tasks are also completed away from school, which contributes to additional sedentary time. Most Australian 4- to 5-year-old children (85%) who are not yet in school attend preschool.²⁷ A recent review found estimates of screen time use during childcare ranges from 0.1 to 2.4 hours per day.²⁸ Sedentary transport tasks include sitting in buses, trains and cars to get to and from school and other destinations. Sedentary self-care tasks include eating and some grooming. Leisure and play sedentary behaviors include reading from a book or an electronic screen. With such a diversity of tasks and differential time spent in each task, it is likely that not all sedentary behaviors are equal in terms of their impact on healthy growth, development, and wellbeing.^{29,30}

Sedentary behaviors are often classified as being either based around an electronic screen or not.^{23,31} Screen time sedentary behaviors were initially TV, then included video games and desktop/laptop computers and now include touch screen tablets and smart phones. Currently data on the use of new touch screen devices by children are very limited, and the development of smart devices has decoupled device and content—children no longer need a TV to watch “TV.” Nonscreen sedentary behaviors of children typically include class time at school, commuting, reading from paper, talking and eating, though with multitasking and the growing integration of technology into daily life, each of these examples could also involve screen time. Figure 3 shows nationally representative Australian data from 2007 and illustrates that total daily sitting time is high from age 9 to 17 years and is composed of around 3.5 hours of screen time and 6 hours of nonscreen time.²⁶ Thus, while screen time is often the focus, it does not constitute the majority of sedentary behavior for most children.

In summary, children spend a large proportion of their waking hours in sedentary behaviors for a range of reasons. Childhood sedentary behavior is varied in aspects potentially important to child health and development and given the high exposure and varied nature of sedentary behavior, it is critical to understand the impact of sedentary behaviors on healthy growth, development, and wellbeing.




Figure 1 — Sedentary behaviors, mechanisms, and impact on child health and development.




Figure 2 — Average proportion of daily wake time spent in “activity” of different intensity for Australian children aged 10 to 12 years (data from Mitchell et al19).

Question 2: What Are the Potential Mechanisms for Sedentary Behaviors to Impact Child Health and Development?

There are a number of mechanisms by which sedentary behaviors may impact child health and development, as illustrated in Figure 1. **Disruption of Metabolism.** Sedentary behaviors could potentially influence energy expenditure, energy intake, and energy metabolism, which could impact adiposity and other cardiometabolic outcomes. Sedentary behaviors may directly decrease energy expenditure. Prolonged low energy expenditure during sedentary behaviors could result in lower daily energy expenditure via low levels of muscle activity and thus decreased energy expenditure. Children typically have low levels of energy expenditure (<1.5 METs) during common sedentary activities.^{10,32} Sedentary behaviors also may displace higher energy expenditure activities, which have clear metabolic health effects. MVPA is known to have positive effects on cardiometabolic outcomes in children including increased myocardial function, improved cholesterol, and decreased blood pressure.^{6,33} Therefore,

children who spend too much time in sedentary behaviors may be in double jeopardy, as they may be impacted by the negative effects of sedentary behaviors and not benefit from the positive effects of the more vigorous activities that could have been engaged in for some of that time.

Some sedentary behaviors, or activities during sedentary behaviors, may directly increase energy intake and thus impact cardiometabolic outcomes. For example, children consumed more calories during a meal while watching TV than while playing with computers or video games.³⁴ Additionally, some sedentary behaviors, or exposure to content during sedentary behaviors, may




Figure 3 — Daily time Australian children spend being sedentary (data from the Australian National Children's Nutrition and Physical Activity Survey²⁵).

indirectly increase later energy intake. For example, increased intake of junk food may result from seeing sugar-sweetened beverage sponsorship signs while watching a sporting event either live or on TV, or viewing fast food advertisements during social media use.^{35,36}

Prolonged sedentary behavior can also alter energy metabolism. Laboratory studies in adults^{37,38} have demonstrated changes in glucose metabolism; however, a similar study in children was not able to demonstrate a similar effect.³⁹ In addition to changes in glucose metabolism, prolonged low energy expenditure may also result in changes in the partitioning of fat and decreased muscle protein synthesis rates⁷ with effects on metabolism occurring beyond time spent in sedentary behaviors. Both the timing and patterns of sedentary behavior may have important influences on energy metabolism.⁴⁰

Limited Neuromuscular Activity. Sedentary behaviors may impact gross motor control and bone and muscle development via low levels of movement and muscle activity and/or the displacement of movement activities with appropriate loading. Lack of practice of gross motor skills could result in reduced motor capacity.⁴¹ Forces exerted during sedentary behavior are typically insufficient to stimulate bone growth, compared with activities such as jumping and skipping.^{42,43} Muscle development similarly requires sufficient loading to stimulate growth, strength development, and flexibility, and sedentary behaviors may not provide sufficient stimulus,⁴⁴ compared with MVPA and strength training.⁴ Some sedentary behaviors may have a positive impact on fine motor skill development, for example, drawing and playing electronic games.⁴⁵

Prolonged, Awkward Postures or Repetitive Motions. Sedentary behaviors could have an impact on musculoskeletal outcomes via prolonged or repetitive stress on tissues. Inflammation of tendons and surrounding connective tissue can be caused by highly repetitive movements, such as video games that require frequent button activation⁴⁵ or playing a piano.⁴⁶ However, these activities may positively impact fine motor skills.⁴⁷ Joint and muscle discomfort can be caused by sustained postures, particularly when the posture is awkward (greater antigravitational load or near to the end of joint range of motion in 1 or more planes), such as writing on paper or watching a video on a smart phone or tablet held close to the body. These activities require positions near to the end range of neck flexion, which may cause neck pain.⁴⁸

Socioemotional Experiences. Sedentary behaviors could have an impact on emotional health and social well-being via exposure to antisocial material and displacement or provision of positive social interaction.⁴⁹ Increased access to the internet adds another avenue for children to be exposed to inappropriate antisocial content and negative social interactions such as cyberbullying.⁵⁰ Sedentary behaviors may also displace or negatively influence useful intrapersonal interactions where children learn social and life skills. Virtual social interactions do not provide all the cues available in face-to-face interactions, and, thus, excessive virtual interaction to the exclusion or even as part of face-to-face interactions may impede a child's social skills.⁵¹ Similarly, other nonsocial nonscreen sedentary behaviors, such as reading books, may have negative developmental psychosocial outcomes.⁵² However, sedentary behaviors such as playing a musical instrument, talking on the phone or video-conferencing with friends and family, and playing multiplayer board and electronic games can provide positive socioemotional experiences.⁵³

Cognitive Experiences. Sedentary behaviors could have an impact on cognitive development and academic achievement by exposure to poor or beneficial cognitive experiences, displacement of more productive sedentary behaviors, and also displacement of MVPA. Some sedentary behaviors encourage passive, rather than active cognitive engagement. Active engagement has shown to have beneficial effects on cognitive development compared with passive activities.⁵⁴ Increased technology use with specific content (eg, content that is hyperstimulating and fast-paced) may have negative effects on children's attention and cognitive performance.⁵⁵ Productive experiences such as school homework may be displaced by other sedentary behavior with limited useful cognitive impact.^{29,30} Additionally, sedentary behaviors displace MVPA, which has been shown to have a positive influence on cognitive performance and academic achievement.⁵⁶ More positively, sedentary behaviors such as appropriate reading, writing, paper and electronic games may have the ability to improve cognitive development and academic achievement.⁵⁷

Other Mechanisms. Sedentary behaviors could have an impact on other aspects of health via a number of mechanisms. Prolonged close vision, for example, reading from a book or tablet, could result in increased short-sightedness.⁵⁸ Sleep quantity and quality could be impacted by bedroom screen time and blue light from some electronic screens altering chrono-hormone levels.⁵⁸

Research supports a link between sedentary behavior and poor health outcomes in adults. One of the pathways by which sedentary behavior may influence health is tracking of the behavior into adulthood. Total sedentary behavior may track better from childhood to adolescence than physical activity.^{59,60} Total screen time behaviors track moderately from childhood to adolescence.⁶¹ TV was more stable than video games from age 5 to 13,⁶² and levels of TV in childhood track into TV in adulthood.⁶³

In summary, there are multiple potential mechanisms for various aspects of sedentary behaviors to impact multiple health and development outcomes. While some mechanisms are specific to certain types of

sedentary behaviors, many may result from a variety of sedentary behaviors. The actual mechanisms are complex, and the interactions and cumulative effects are not fully understood. However, given the considerable exposure of children to sedentary behaviors, it is critical that these relationships are better understood.

Question 3: What Are the Effects of Different Types of Sedentary Behaviors on Child Health and Development?

Sedentary behavior in children has the potential to influence health and development through different types of sedentary behavior and different mechanisms as seen in Figure 1. This section provides a brief synthesis of the available evidence for different sedentary behaviors to have effects on multiple components of child health and development including cardiometabolic, neuromusculoskeletal, psychosocial, and other relevant outcomes. The focus of this brief review is on children, though where the evidence for children is limited,⁶⁴ evidence in adults has been included.⁷ Given the differences in types of sedentary behavior, this brief synthesis is arranged by types of sedentary behavior and includes screen time, TV, other screens (excluding TV), nonscreen sedentary behavior, and any sedentary time.

Screen Time Sedentary Behaviors. The Australian Physical Activity Report Card grades were assigned based on compliance with screen time guidelines, as screen time has been given particular attention for having unique effects on children's health.⁴⁹ Common limitations to the evidence, however, include cross-sectional designs and that many of the observed associations have a high risk of residual confounding due to sedentary behaviors being related to other lifestyle and socioeconomic factors.

Cardiometabolic: The 2 most commonly studied cardiometabolic outcomes have been obesity and cardiorespiratory fitness. A longitudinal study of Danes found that increased TV and total screen time from adolescence to adulthood was associated with increased body mass index (BMI).⁶⁵ A cross-sectional study of 9- to 16-year-olds found that BMI was more strongly inversely associated with general screen time than physical activity.⁶⁶ Cross-sectional studies have also shown a negative relationship between screen time and cardiorespiratory fitness that is independent of physical activity.^{67,68}

Neuromusculoskeletal: The majority of studies examining musculoskeletal effects of screen time has examined specific types of screens and will thus be discussed in following sections. However, in 1 cross-sectional study, overall screen time was not associated with bone structure in 9- to 20-year-old children when adjusted for physical activity and other factors.⁴²

Psychosocial: Compared with other types of sedentary behavior, screen time has a unique potential to influence psychosocial outcomes due to the content viewed. While the assumption is that screen time negatively affects psychosocial outcomes, few studies have empirically evaluated this relationship. Two cross-sectional studies found increased screen time to be detrimentally associated with depression scores and psychological difficulty, independent of physical activity.^{69,70} Additionally, evidence supports the transmission of aggressive behaviors from violent media including TV, movies, video games, and the Internet.⁷¹ Specific uses of technology, such as for educational purposes, can, nevertheless, improve psychosocial outcomes, and these are discussed in later sections.

Other: Unique characteristics of screen time behavior have also led to the investigation of other outcomes from screen time including sleep and vision. Among adults, screen time, not total sedentary time, was associated with sleep problems.⁷² A review found that increased screen time among children adversely affected sleep, but the effects largely depended on type of screen exposure, age, gender, and day of the week.⁷³ Screen time may also adversely affect vision. Among university students, sustained periods of

close screen work and lack of a screen filter was associated with a greater report of vision problems including dry and tired eyes as well as headache.⁷⁴

Television Watching. While many of the Physical Activity Report Cards assessed children's exposure to sedentary behaviors based on meeting guidelines for total screen time, it is acknowledged that different types of screen devices, used for different purposes, may have differential effects on child health and development. The majority of the evidence supports a detrimental effect of TV on multiple child outcomes.

Cardiometabolic: Several cross-sectional studies support an inverse relationship between TV and cardiometabolic risk in children independent of physical activity.^{75–78} These studies have varied in age group and how they have accounted for physical activity.

Additional cross-sectional studies have examined the relationship between TV and BMI, but few studies have tested causal relationships. In a worldwide study of children aged 5 to 15 years, there was a positive association between TV and BMI, but the relationship was not adjusted for physical activity.⁷⁹ In a longitudinal study in The Netherlands, an increase in TV from adolescence to adulthood was associated with increased BMI in adulthood.⁶⁵

There is a lack of evidence to support a relationship between TV and cardiorespiratory fitness in children. A longitudinal study found that increased TV was associated with decreased cardiorespiratory fitness over 2 years from age 7, but this was not adjusted for physical activity.⁸⁰ In female adults, TV was negatively associated with cardiorespiratory fitness, but this was mostly mediated by PA and percent body fat.⁸¹

Independent of total sedentary and screen time, TV may have additional harmful effects on energy balance due to its relationship with energy intake. Several cross-sectional studies have found an association between increased TV and a poorer diet.^{82–84} An experimental study found that energy intake increased while watching TV among 9- to 13-year-olds.³⁴ Advertising during TV may also lead to subsequent increased energy intake, as shown in experimental studies.^{35,36}

Neuromusculoskeletal: The evidence for the effects of TV on neuromusculoskeletal outcomes in children is inconclusive. While 1 study has found that TV and back pain were positively related,⁸⁵ 2 others have found that TV was not related to back pain⁸⁶ or neck and back pain.⁸⁷

Psychosocial: A large number of studies have examined relationships between TV and various psychosocial effects; however, many of them have been cross sectional and unable to discern causality. The majority have found negative associations between increased TV and psychosocial outcomes. Research suggests that children who watch more TV are more likely to have behavioral difficulties, but a variety of measures and definitions of behavior have been used.^{70,88,89} In a longitudinal study of preschoolers aged 2 to 3 years, TV was positively associated with externalizing problems.⁹⁰ Other psychological outcomes have been found to have crosssectional associations with TV, without adjustment for physical activity, including psychological distress,⁹¹ self-esteem,⁹² criminal conviction, antisocial personality disorder, and aggressive traits.⁹³ While an association between TV and aggressive behavior has been suggested, the evidence is unclear.⁹⁴ Cross-sectional associations suggest that children who watch more TV have poorer cognitive performance including executive function,⁹⁵ communication and language development,⁹⁶ and hyperactivity/inattention.⁹⁷

Other: Both vision and sleep seem to be negatively affected by increased TV. TV (and computer use) was associated with poorer vision in children aged 6 to 18 years.⁹⁸ Increased TV has been associated with poorer sleep in 2 longitudinal studies, including shorter sleep time unadjusted for physical activity in a

longitudinal study of children from 6 months to 7 years⁹⁹ and from ages 2 to 4 and 6 to 9 when adjusted for parent-reported PA.¹⁰⁰

Other Screens (Not TV). There have been few studies to isolate other screens (not including TV), with most of them examining computer use or electronic video games.

Cardiometabolic: Saunders et al found that leisure-time computer/video game play in boys (TV in girls) was associated with poorer cardiometabolic profiles among 8- to 11-year-olds when adjusted for accelerometer determined physical activity.⁷⁷ Another cross-sectional study reported computer game use was positively associated with overweight status in 6- to 14-year-old children but not in highly active children.¹⁰¹

Neuromusculoskeletal: The associations between technology and low back and neck/shoulder pain have been inconsistent. Crosssectional surveys of adolescents have found computer and laptop use, greater than 2 hours, were associated with low back and neck/ shoulder pain.^{87,102} However, another cross-sectional study of adolescents found that neck/shoulder pain was not related to computer use when adjusted for physical activity.¹⁰³ Among children, neck pain was related to increased computer use⁵⁸ and repetitive electronic game use has been shown to be related to tendonitis.⁴⁵ However, cross-sectional evidence suggests that young children who play greater amounts of interactive video games have improved object control motor skills.¹⁰⁴

Psychosocial: Numerous studies have examined the relationship between other screens, particularly computers and video games, with both positive and negative psychosocial outcomes. The majority have been cross sectional, which again limits the ability to support causal relationships. A meta-analysis found that violent video game play was related to increased aggressive behavior, aggressive cognition, and aggressive affect and decreased empathy and prosocial behavior.¹⁰⁵ Time playing video games has been cross-sectionally related to negative outcomes such as depression, lower academic achievement, conduct problems,¹⁰⁶ and poorer working memory,¹⁰⁷ whereas high amounts of computer use have been associated with weaker performance in tests measuring flexibility of attention.¹⁰⁷

While many of the studies have found detrimental associations, there is also evidence for benefits of other types of screen use. A cross-sectional study of adolescents found that self-reported video usage was positively correlated with improvements in brain structures that correlate with improved executive function.¹⁰⁸ In educational research, technology use (laptops and tablets) has been shown to improve educational outcomes, but often the study designs were weak with small samples and no comparison groups.¹⁰⁹ Technology may be especially beneficial for those with learning disabilities.^{110,111} Despite concerns over children becoming technology dependent and losing social interaction skills, adolescents who had more smartphone use also had more face-to-face interactions.¹¹²

Other: Computer use has been cross-sectionally associated with poorer vision in 6- to 18-year-old children.⁹⁸ Other media use, compared with TV, was more strongly correlated to health and wellbeing among 8- to 13-year-olds, though this was not adjusted for physical activity.¹¹³

Nonscreen Sedentary Behaviors. Nonscreen sedentary behaviors have also been related to various health and development outcomes, but the heterogeneity of behaviors and outcomes precludes a comprehensive review in this paper. Further, much of the research has not separated nonscreen sedentary behavior from other sedentary behaviors. A few examples are, nevertheless, provided to illustrate how nonscreen sedentary behaviors may influence health. Puzzle play in early childhood has been associated with improved spatial abilities.¹¹⁴ Unsurprisingly, increased time spent reading during school was related

to higher reading achievement, although time spent reading at home was not.¹¹⁵ Sedentary practices such as meditation are associated with improved cognitive process¹¹⁶ and self-esteem in school children.¹¹⁷

Total Sedentary Time. *Cardiometabolic:* Total sedentary time, in activities with a low energy expenditure, has been associated with several cardiometabolic outcomes in a recent review,⁶ although, after adjusting for MVPA, the evidence was inconsistent.¹⁶ The strength of association depends on the specific variables examined. For example, in a cross-sectional study of multiple cardiometabolic outcomes among 5- to 10-year-old children, only high-density lipoprotein cholesterol was negatively associated with sedentary time measured by accelerometry, independent of physical activity.¹⁷

BMI has been the most common cardiometabolic outcome measured, yet even the evidence for this relationship has been inconsistent. In adults, a positive relationship between sedentary time and BMI has been found, independent of physical activity.¹¹⁸ However, a recent review of longitudinal studies among children has concluded that the evidence to support a relationship between sedentary behavior and adiposity is inconclusive.¹¹⁹ Reasons for the inconclusive findings may be the predominance of cross-sectional studies, varying measures of sedentary time, and inconsistent adjustment for physical activity.³¹ One problem with measuring sedentary time with accelerometers may be the misclassification of standing time as sedentary.¹²⁰

Similar to BMI and adiposity, the relationship between sedentary time and cardiorespiratory fitness has been inconsistent. In adults, a large cross-sectional study using National Health and Nutrition Examination Survey (NHANES) data found an inverse association between total sedentary time and cardiorespiratory fitness, even when adjusted for exercise.¹³ Comparatively in children, a cross-sectional study of over 2,000 10- to 18-year-olds did not find an independent relationship between cardiorespiratory fitness and total sedentary time when also adjusted for physical activity.¹⁸ Additional evidence suggests that the relationship may differ between genders.¹²¹

Of particular interest to cardiometabolic outcomes may be sedentary time accumulated in long, uninterrupted bouts. Literature in adults suggests that these long, uninterrupted bouts may be particularly detrimental,^{14,122} though the evidence in children has been less conclusive and predominantly cross sectional.^{17,77,123} In 1 randomized crossover study, breaking up long bouts of sedentary behavior in 10- to 14-year-olds did not result in changes to cardiometabolic markers.³⁹

Neuromusculoskeletal: Few studies have examined the relationship between total sedentary time and neuromusculoskeletal outcomes including motor skills, bone structure, and musculoskeletal discomfort or pain. One cross-sectional study found that increased sedentary time was negatively associated with motor proficiency among 9- to 10-year-olds, independent of physical activity.¹²⁴ Another cross-sectional study examined bone structure and found no association with total sedentary time when adjusted for physical activity.⁴² Finally, there has been inconsistent evidence for sedentary time to be related to musculoskeletal pain in children.¹²⁵⁻¹²⁷

Psychosocial: Of the multiple psychosocial outcomes that may be potentially affected by sedentary time, very few studies have studied relationships with sedentary time. Two cross-sectional studies have found no associations with self-esteem,¹²⁸ and negative associations with sustained attention but no other tests in a cognitive battery.¹⁰⁷

Other: Total sedentary time may also be associated with other health related outcomes. In adults, there is an increased risk of allcause mortality with daily sitting time greater than 8 hours per day independent of physical activity.¹²⁹

In summary, there is considerable evidence showing sedentary behaviors have implications for child health and development. However, the strength of current evidence varies by types of sedentary behavior and health outcomes, as well as the methodological approaches used to examine these relationships.

Further Research Needed to Inform Strategies to Improve the Grade To better understand which sedentary behaviors are occurring and answer Question 1, national surveillance systems are required to provide robust estimates of children's sedentary behavior exposure. Data are required from infancy, across childhood to adulthood and need to examine the different types of sedentary behaviors, the different devices used while sedentary, the content or tasks performed and the context of behavior. Data should also be tracked longitudinally.

To better understand the mechanisms for these impacts and answer Question 2, mechanistic studies are required to test causal pathways and inform critical components for interventions. To better understand the impact of these behaviors and answer Question 3, longitudinal and experimental design studies are required to provide stronger causal evidence of the impacts of the various sedentary behaviors on the full range of important child health and developmental outcomes. Analyses need to consider dose-response relationships while also evaluating mediating and moderating influences such as physical activity, built environment, family socioeconomic status and parenting style. More sophisticated statistical approaches are needed, for example compositional analysis may be useful when considering the limited 24-hour nature of each day, which can be divided into exhaustive and mutually exclusive components.¹³⁰ A life-course approach can be used to evaluate critical windows and pathways of causality.

Further research is needed to improve the measurement of both the amount and nature of children's sedentary behaviors and which strategies are effective to improve sedentary behaviors. Sedentary behavior measurement needs to be improved to encompass a whole-of-day approach, including sleep and wake time and the full spectrum of wake time "activity." Measurement needs to capture not just the total amount of exposure, but also the pattern of exposure and the potential overlap of behaviors with multitasking. Methods to accurately capture the context and content/task/device details of behaviors also need to be developed. Standardized and practical methods for classifying and quantifying sedentary behaviors need to be developed to enable valid comparisons between countries. These methods need to match understandings of mechanisms and, thus, key aspects of behavior to capture. For example, using inclinometers to measure total sedentary time or validated technology monitoring apps to measure content, accumulation and pattern of screen time. Reevaluation and refinement of partitioning of "activity" into different intensity-based categories also needs to be conducted, to understand the postural or energy expenditure aspects which relate to outcomes. Comparisons should also be undertaken of countries with healthier sedentary exposure for their children to determine whether some aspects of that society can be promoted in countries with poorer sedentary behavior grades.

Finally, while not reviewed in this paper, continued intervention research is needed to evaluate the efficacy (do the interventions produce a desired effect) and cost efficiency (are the interventions economical) of various strategies to improve sedentary behavior exposure in children.¹³¹ Reviews of studies evaluating various strategies would provide useful guidance on policies and interventions to be promoted. The importance of tailoring interventions to specific groups of children (age group, gender, socioeconomic status, leisure interests, etc.) and targeting specific behavior change (video games, book reading, passive transport, etc.) also needs to be evaluated.

Conclusion

The available evidence, while incomplete,⁶⁴ is sufficiently convincing that sedentary behaviors are critical to child health and development. Nations therefore need to have strategies to promote appropriate

exposure to these common behaviors. It appears likely that both the total exposure and pattern of exposure are important for cardiometabolic and neuromusculoskeletal outcomes and so there is a need to reduce overall sedentary time and prolonged bouts of sedentary time for many children. Aspects of sedentary tasks, such as content, device and context, also appear important to a range of outcomes including psychosocial outcomes and thus need to be addressed.

Failure to adequately address this issue is likely to result in nations facing unsustainable health and economic burdens for poor child and adult health and developmental outcomes. A range of intervention options is available in all nations, targeting the child directly or indirectly via parents, teachers/schools, peers, technology, and societal infrastructure. Nations can therefore look forward to improving their grade based on the sedentary behavior of their children, if they invest sufficiently in understanding this key behavior and in strategies to promote appropriate behavior.

Acknowledgments

Straker is supported by a National Health and Medical Research Council of Australia Senior Research Fellowship (1019980). Cliff is funded by an Australian Research Council Discovery Early Career Researcher Award (DE140101588). Gomersall is supported by a National Health and Medical Research Council Program Grant (569940). The Australian Report Card on Physical Activity for Children and Young People Research Working Group also included Billie Giles-Corti, Louise Hardy, Kylie Hesketh, Andrew Hills, David Lubans, Cuane Macdonald, Rona Macniven, Philip Morgan, Tony Okley, Anne-Maree Parish, Ron Plotnikoff, Trevor Shilton, Anna Timperio, Stewart Trost, and Stewart Vella.

References

1. Department of Health. *Australia's Physical Activity and Sedentary Behaviour Guidelines*. Canberra, Australia: Commonwealth of Australia. 2014.
2. Schranz N, Olds T, Cliff D, et al. Results from Australia's 2014 report card on physical activity for children and youth. *J Phys Act Health*. 2014;11(4 Suppl 1):S21–S25. PubMed doi:10.1123/jpah.2014-0164
3. Australian Bureau of Statistics. *Australian Health Survey: Physical Activity, 2011–12*. Catalogue No. 4364.0. Canberra, Australia: Author; 2013.
4. Okely AD, Salmon J, Vella SA, et al. *A Systematic Review to inform the Australian Sedentary Behaviour Guidelines for Children and Young People*. Canberra, Australia: Commonwealth of Australia; 2012.
5. Tremblay MS, LeBlanc AG, Kho ME, et al. Systematic review of sedentary behaviour and health indicators in school-aged children and youth. *Int J Behav Nutr Phys Act*. 2011;8:98. PubMed doi:10.1186/1479-5868-8-98
6. Saunders TJ, Chaput JP, Tremblay MS. Sedentary behaviour as an emerging risk factor for cardiometabolic diseases in children and youth. *Can J Diabetes*. 2014;38(1):53–61. PubMed doi:10.1016/j.jcjd.2013.08.266
7. Thyfault JP, Du M, Kraus WE, Levine JA, Booth FW. Physiology of sedentary behavior and its relationship to health outcomes. *Med Sci Sports Exerc*. 2015;47(6):1301–1305. PubMed doi:10.1249/mss.0000000000000518

8. Sedentary Behaviour Research Network. Standardized use of the terms “sedentary” and “sedentary behaviours.” *Appl Physiol Nutr Metab*. 2012;37:540–542. PubMed doi:10.1139/h2012-024
9. Cain KL, Sallis JF, Conway TL, Van Dyck D, Calhoun L. Using accelerometers in youth physical activity studies: a review of methods. *J Phys Act Health*. 2013;10(3):437–450.
10. Reilly JJ, Janssen X, Cliff DP, Okely AD. Appropriateness of the definition of “sedentary” in young children: whole-room calorimetry study. *J Sci Med Sport*. 2015;18(5):565–568. PubMed doi: 10.1016/j.jsams.2014.07.013
11. Lubans DR, Hesketh K, Cliff DP, et al. A systematic review of the validity and reliability of sedentary behaviour measures used with children and adolescents. *Obes Rev*. 2011;12(10):781–799. PubMed doi:10.1111/j.1467-789X.2011.00896.x
12. Pate RR, Mitchell JA, Byun W, Dowda M. Sedentary behaviour in youth. *Br J Sports Med*. 2011;45(11):906–913. PubMed doi:10.1136/bjsports-2011-090192
13. Kulinski JP, Khera A, Ayers CR, et al. Association between cardiorespiratory fitness and accelerometer-derived physical activity and sedentary time in the general population. *Mayo Clin Proc*. 2014;89(8):1063–1071. PubMed doi:10.1016/j.mayocp.2014.04.01914.
14. Latouche C, Jowett JB, Carey AL, et al. Effects of breaking up prolonged sitting on skeletal muscle gene expression. *J Appl Physiol*. 2013;114(4):453–460.
15. Dunstan DW, Thorp AA, Healy GN. Prolonged sitting: is it a distinct coronary heart disease risk factor? *Curr Opin Cardiol*. 2011;26(5):412–419. PubMed doi:10.1097/HCO.0b013e3283496605
16. Ekelund U, Luan J, Sherar LB, Esliger DW, Griew P, Cooper A. Moderate to vigorous physical activity and sedentary time and cardiometabolic risk factors in children and adolescents. *JAMA*. 2012;307(7):704–712. PubMed doi:10.1001/jama.2012.156
17. Cliff DP, Jones RA, Burrows TL, et al. Volumes and bouts of sedentary behavior and physical activity: associations with cardiometabolic health in obese children. *Obesity*. 2014;22(5):E112–E118. PubMed doi:10.1002/oby.20698
18. Marques A, Santos R, Ekelund U, Sardinha LB. Association between physical activity, sedentary time and healthy fitness in youth. *Med Sci Sports Exerc*. 2015;47(3):575–580. PubMed doi:10.1249/mss.0000000000000426
19. Mitchell JA, Pate RR, Beets MW, Nader PR. Time spent in sedentary behavior and changes in childhood BMI: a longitudinal study from ages 9 to 15 years. *Int J Obes*. 2013;37(1):54–60. PubMed doi:10.1038/ijo.2012.41
20. Abbott RA, Straker LM, Mathiassen SE. Patterning of children’s sedentary time at and away from school. *Obesity*. 2013;21(1):E131–E133. PubMed doi:10.1002/oby.20127
21. Hnatiuk J, Ridgers ND, Salmon J, Campbell K, McCallum Z, Hesketh K. Physical activity levels and patterns of 19-month-old children. *Med Sci Sports Exerc*. 2012;44(9):1715–1720. PubMed doi:10.1249/MSS.0b013e31825825c4
22. Hinkley T, Salmon J, Okely AD, Crawford D, Hesketh K. Preschoolers’ physical activity, screen time, and compliance with recommendations. *Med Sci Sports Exerc*. 2012;44(3):458–465. PubMed doi:10.1249/MSS.0b013e318233763b

23. Straker L, Smith A, Hands B, Olds T, Abbott R. Screen-based media use clusters are related to other activity behaviours and health indicators in adolescents. *BMC Public Health*. 2013;13:1174. PubMed doi:10.1186/1471-2458-13-1174
24. Carson V, Cliff DP, Janssen X, Okely AD. Longitudinal levels and bouts of sedentary time among adolescent girls. *BMC Pediatr*. 2013;13:173. PubMed doi:10.1186/1471-2431-13-173
25. Currie C, Zanotti C, Morgan A, et al, eds. *Social determinants of health and well-being among young people: Health Behaviour in School-aged Children (HBSC) study: international report from the 2009/2010 survey (Health Policy for Children and Adolescents, No. 6)*. Copenhagen, Denmark: World Health Organization Regional Office for Europe; 2012.
26. Department of Health and Ageing Australian National Children's Nutrition and Physical Activity Survey: main findings. Report prepared by Commonwealth Scientific and Industrial Research Organisation (CSIRO) and the University of South Australia. Canberra, Australia: Department of Health and Ageing. 2008
27. Australian Bureau of Statistics. *Childhood Education and Care*. Australia. 2011.
28. Vanderloo LM. Screen-viewing among preschoolers in childcare: a systematic review. *BMC Pediatr*. 2014;14:205. PubMed doi:10.1186/1471-2431-14-205
29. Christakis DA. Interactive media use at younger than the age of 2 years: time to rethink the American Academy of Pediatrics guideline? *JAMA Pediatr*. 2014;168(5):399–400. PubMed doi:10.1001/jamapediatrics.2013.5081
30. Voss MW, Carr LJ, Clark R, Weng T. Revenge of the “sit” II: does lifestyle impact neuronal and cognitive health through distinct mechanisms associated with sedentary behavior and physical activity? *Ment Health Phys Act*. 2014;7(1):9–24. doi:10.1016/j.mhpa.2014.01.001
31. Foley LS, Maddison R, Jiang Y, Olds T, Ridley K. It's not just the television: survey analysis of sedentary behaviour in New Zealand young people. *Int J Behav Nutr Phys Act*. 2011;8:132. PubMed doi:10.1186/1479-5868-8-132
32. Straker L, Abbott R. Effect of screen based media on energy expenditure and heart rate in 9–12-year-old children. *Pediatr Exerc Sci*. 2007;19:459–471. PubMed
33. Janssen I, LeBlanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act*. 2010;7:40. PubMed
34. Marsh S, Ni Mhurchu C, Jiang Y, Maddison R. Comparative effects of TV watching, recreational computer use, and sedentary video game play on spontaneous energy intake in male children: a randomised crossover trial. *Appetite*. 2014;77:13–18. PubMed doi:10.1016/j.appet.2014.02.008
35. Boyland EJ, Halford JC. Television advertising and branding: effects on eating behaviour and food preferences in children. *Appetite*. 2013;62:236–241. PubMed doi:10.1016/j.appet.2012.01.032
36. Boyland EJ, Harrold JA, Kirkham TC, et al. Food commercials increase preference for energy-dense foods, particularly in children who watch more television. *Pediatrics*. 2011;128(1):e93–e100. PubMed doi:10.1542/peds.2010-1859
37. Dunstan DW, Kingwell BA, Larsen R, et al. Breaking up prolonged sitting reduces postprandial glucose and insulin responses. *Diabetes Care*. 2012;35(5):976–983. PubMed doi:10.2337/dc11-1931

38. Peddie MC, Bone JL, Rehner NJ, Skeaff CM, Gray AR, Perry TL. Breaking prolonged sitting reduces postprandial glycemia in healthy, normal-weight adults: a randomized crossover trial. *Am J Clin Nutr*. 2013;98(2):358–366. doi:10.3945/ajcn.112.051763
39. Saunders TJ, Chaput JP, Goldfield GS, et al. Prolonged sitting and markers of cardiometabolic disease risk in children and youth: a randomized crossover study. *Metabolism*. 2013;62(10):1423–1428. PubMed doi:10.1016/j.metabol.2013.05.010
40. Golley RK, Maher CA, Matricciani L, Olds TS. Sleep duration or bedtime? Exploring the association between sleep timing behaviour, diet and BMI in children and adolescents. *Int J Obes*. 2013;37(4):546–551. PubMed doi:10.1038/ijo.2012.212
41. Straker LM, Campbell AC, Jensen LM, et al. Rationale, design and methods for a randomised and controlled trial of the impact of virtual reality games on motor competence, physical activity, and mental health in children with developmental coordination disorder. *BMC Public Health*. 2011;11:654. PubMed doi:10.1186/1471-2458-11-654
42. Gabel L, McKay HA, Nettlefold L, Race D, Macdonald HM. Bone architecture and strength in the growing skeleton: the role of sedentary time. *Med Sci Sports Exerc*. 2015;47(2):363–372. PubMed doi:10.1249/mss.0000000000000418
43. McKay H, Liu D, Egeli D, Boyd S, Burrows M. Physical activity positively predicts bone architecture and bone strength in adolescent males and females. *Acta Paediatr*. 2011;100(1):97–101. PubMed doi:10.1111/j.1651-2227.2010.01995.x
44. Straker L, Mathiassen SE. Increased physical work loads in modern work: a necessity for better health and performance? *Ergonomics*. 2009;52(10):1215–1225. PubMed doi:10.1080/00140130903039101
45. Straker L, Abbott R, Collins R, Campbell A. Evidence-based guidelines for wise use of electronic games by children. *Ergonomics*. 2014;57(4):471–489. PubMed doi:10.1080/00140139.2014.895856
46. Ranelli S, Straker L, Smith A. Playing-related musculoskeletal problems in children learning instrumental music: the association between problem location and gender, age, and music exposure factors. *Med Probl Perform Art*. 2011;26(3):123–139. PubMed
47. Costa-Giomi E. Does music instruction improve fine motor abilities? *Ann N Y Acad Sci*. 2005;1060(1):262–264. PubMed doi:10.1196/annals.1360.053
48. Straker LM, Coleman J, Skoss R, Maslen BA, Burgess-Limerick R, Pollock CM. A comparison of posture and muscle activity during tablet computer, desktop computer and paper use by young children. *Ergonomics*. 2008;51(4):540–555. PubMed doi:10.1080/00140130701711000
49. Sigman A. Time for a view on screen time. *Arch Dis Child*. 2012;97(11):935–942. PubMed doi:10.1136/archdischild-2012-302196
50. Modecki KL, Minchin J, Harbaugh AG, Guerra NG, Runions KC. Bullying prevalence across contexts: a meta-analysis measuring cyber and traditional bullying. *J Adolesc Health*. 2014;55(5):602–611. PubMed doi:10.1016/j.jadohealth.2014.06.007
51. Przybylski AK, Weinstein N. Can you connect with me now? How the presence of mobile communication technology influences face-to-face conversation quality. *J Soc Pers Relat*. 2013;30(3):237–246. doi:10.1177/0265407512453827

52. McHale SM, Crouter AC, Tucker CJ. Free-time activities in middle childhood: links with adjustment in early adolescence. *Child Dev.* 2001;72(6):1764–1778. PubMed doi:10.1111/1467-8624.00377
53. Ryan RM, Rigby CS, Przybylski A. The motivational pull of video games: a self-determination theory approach. *Motiv Emot.* 2006;30(4):344–360. doi:10.1007/s11031-006-9051-8
54. Corno L, Mandinach EB. The role of cognitive engagement in classroom learning and motivation. *Educ Psychol.* 1983;18(2):88–108. doi:10.1080/00461528309529266
55. Lillard AS, Peterson J. The immediate impact of different types of television on young children's executive function. *Pediatrics.* 2011;128(4):644–649. PubMed doi:10.1542/peds.2010-1919
56. Biddle SJ, Asare M. Physical activity and mental health in children and adolescents: a review of reviews. *Br J Sports Med.* 2011;45(11):886–895. PubMed doi:10.1136/bjsports-2011-090185
57. Haapala EA, Poikkeus AM, Kukkonen-Harjula K, et al. Associations of physical activity and sedentary behavior with academic skills: a follow-up study among primary school children. *PLoS One.* 2014;9(9):e107031. PubMed doi:10.1371/journal.pone.0107031
58. Straker L, Pollock C, Maslen B. Principles for the wise use of computers by children. *Ergonomics.* 2009;52(11):1386–1401. PubMed doi:10.1080/00140130903067789
59. Biddle SJ, Pearson N, Ross GM, Braithwaite R. Tracking of sedentary behaviours of young people: a systematic review. *Prev Med.* 2010;51(5):345–351. PubMed doi:10.1016/j.ypmed.2010.07.018
60. Jones RA, Hinkley T, Okely AD, Salmon J. Tracking physical activity and sedentary behavior in childhood: a systematic review. *Am J Prev Med.* 2013;44(6):651–658. PubMed doi:10.1016/j.amepre.2013.03.001
61. Gebremariam MK, Totland TH, Andersen LF, et al. Stability and change in screen-based sedentary behaviours and associated factors among Norwegian children in the transition between childhood and adolescence. *BMC Public Health.* 2012;12:104. PubMed doi:10.1186/1471-2458-12-104
62. Francis SL, Stancel MJ, Sernulka-George FD, Broffitt B, Levy SM, Janz KF. Tracking of TV and video gaming during childhood: Iowa Bone Development Study. *Int J Behav Nutr Phys Act.* 2011;8:100. PubMed doi:10.1186/1479-5868-8-100
63. Smith L, Gardner B, Hamer M. Childhood correlates of adult TV viewing time: a 32-year follow-up of the 1970 British Cohort Study. *J Epidemiol Community Health.* 2015;69(4):309–313. PubMed doi:10.1136/jech-2014-204365
64. de Rezende LF, Rodrigues Lopes M, Rey-Lopez JP, Matsudo VK, Luiz Odo C. Sedentary behavior and health outcomes: an overview of systematic reviews. *PLoS One.* 2014;9(8):e105620. PubMed doi:10.1371/journal.pone.0105620
65. Grøntved A, Ried-Larsen M, Møller NC, et al. Youth screen-time behaviour is associated with cardiovascular risk in young adulthood: the European Youth Heart Study. *Eur J Prev Cardiol.* 2014;21(1):49–56. PubMed doi:10.1177/2047487312454760
66. Maher C, Olds TS, Eisenmann JC, Dollman J. Screen time is more strongly associated than physical activity with overweight and obesity in 9- to 16-year-old Australians. *Acta Paediatr.* 2012;101(11):1170–1174. PubMed doi:10.1111/j.1651-2227.2012.02804.x

67. Mitchell JA, Pate RR, Blair SN. Screen-based sedentary behavior and cardiorespiratory fitness from age 11 to 13. *Med Sci Sports Exerc.* 2012;44(7):1302–1309. PubMed doi:10.1249/MSS.0b013e318247cd73
68. Sandercock GR, Ogunleye AA. Screen time and passive school travel as independent predictors of cardiorespiratory fitness in youth. *Prev Med.* 2012;54(5):319–322. PubMed doi:10.1016/j.ypmed.2012.03.007
69. Kremer P, Elshaug C, Leslie E, Toumbourou JW, Patton GC, Williams J. Physical activity, leisure-time screen use and depression among children and young adolescents. *J Sci Med Sport.* 2014;17(2):183–187. PubMed doi:10.1016/j.jsams.2013.03.012
70. Page AS, Cooper AR, Griew P, Jago R. Children's screen viewing is related to psychological difficulties irrespective of physical activity. *Pediatrics.* 2010;126(5):e1011–e1017. PubMed doi:10.1542/peds.2010-1154
71. Browne KD, Hamilton-Giachritsis C. The influence of violent media on children and adolescents: a public-health approach. *Lancet.* 2005;365(9460):702–710.
72. Vallance JK, Buman MP, Stevinson C, Lynch BM. Associations of overall sedentary time and screen time with sleep outcomes. *Am J Health Behav.* 2015;39(1):62–67. PubMed doi:10.5993/AJHB.39.1.7
73. Hale L, Guan S. Screen time and sleep among school-aged children and adolescents: a systematic literature review. *Sleep Med Rev.* 2015;21:50–58. PubMed doi:10.1016/j.smrv.2014.07.007
74. Shantakumari N, Eldeeb R, Sreedharan J, Gopal K. Computer use and vision-related problems among university students in Ajman, United Arab Emirate. *Ann Med Health Sci Res.* 2014;4(2):258–263. PubMed doi:10.4103/2141-9248.129058
75. Stamatakis E, Coombs N, Jago R, et al. Type-specific screen time associations with cardiovascular risk markers in children. *Am J Prev Med.* 2013;44(5):481–488. PubMed doi:10.1016/j.amepre.2013.01.020
76. Väistö J, Eloranta AM, Viitasalo A, et al. Physical activity and sedentary behaviour in relation to cardiometabolic risk in children: cross-sectional findings from the Physical Activity and Nutrition in Children (PANIC) Study. *Int J Behav Nutr Phys Act.* 2014;11:55. PubMed doi:10.1186/1479-5868-11-55
77. Saunders TJ, Tremblay MS, Mathieu ME, et al. Associations of sedentary behavior, sedentary bouts and breaks in sedentary time with cardiometabolic risk in children with a family history of obesity. *PLoS One.* 2013;8(11):e79143. PubMed doi:10.1371/journal.pone.0079143
78. Staiano AE, Abraham AA, Calvert SL. Adolescent exergame play for weight loss and psychosocial improvement: a controlled physical activity intervention. *Obesity.* 2013;21(3):598–601. PubMed doi:10.1002/oby.20282
79. Braithwaite I, Stewart AW, Hancox RJ, Beasley R, Murphy R, Mitchell EA. The worldwide association between television viewing and obesity in children and adolescents: cross sectional study. *PLoS One.* 2013;8(9):e74263. PubMed doi:10.1371/journal.pone.0074263
80. Mota J, Ribeiro JC, Carvalho J, Santos MP, Martins J. Television viewing and changes in body mass index and cardiorespiratory fitness over a 2-year period in schoolchildren. *Pediatr Exerc Sci.* 2010;22(2):245–253. PubMed

81. Tucker LA, Arens PJ, Lecheminant JD, Bailey BW. Television viewing time and measured cardiorespiratory fitness in adult women. *Am J Health Promot.* 2015;29(5):285–290. PubMed doi:10.4278/ajhp.131107-QUAN-565
82. Olafsdottir S, Eiben G, Prell H, et al. Young children's screen habits are associated with consumption of sweetened beverages independently of parental norms. *Int J Public Health.* 2014;59(1):67–75. PubMed doi:10.1007/s00038-013-0473-2
83. Borghese MM, Tremblay MS, Leduc G, et al. Independent and combined associations of total sedentary time and television viewing time with food intake patterns of 9- to 11-year-old Canadian children. *Appl Physiol Nutr Metab.* 2014;39(8):937–941. PubMed
84. Sisson SB, Shay CM, Broyles ST, Leyva M. Television-viewing time and dietary quality among U.S. children and adults. *Am J Prev Med.* 2012;43(2):196–200. PubMed doi:10.1016/j.amepre.2012.04.016
85. Balagué F, Troussier B, Salminen JJ. Non-specific low back pain in children and adolescents: risk factors. *Eur Spine J.* 1999;8(6):429–438. PubMed doi:10.1007/s005860050201
86. Gunzburg R, Balague F, Nordin M, et al. Low back pain in a population of school children. *Eur Spine J.* 1999;8(6):439–443. PubMed doi:10.1007/s00586005020287.
87. Hakala PT, Rimpela A, Saarni L, Salminen J. Frequency computerrelated activities increase the risk of neck-shoulder and low back pain in adolescents. *Eur J Public Health.* 2006;16(5):536–541. PubMed doi:10.1093/eurpub/ckl025
88. Yousef S, Eapen V, Zoubeidi T, Mabrouk A. Behavioral correlation with television watching and videogame playing among children in the United Arab Emirates. *Int J Psychiatry Clin Pract.* 2014;18(3):203–207. PubMed doi:10.3109/13651501.2013.874442
89. Parkes A, Sweeting H, Wight D, Henderson M. Do television and electronic games predict children's psychosocial adjustment? Longitudinal research using the U.K. Millennium Cohort Study. *Arch Dis Child.* 2013;98(5):341–348. PubMed doi:10.1136/archdischild-2011-301508
90. Verlinden M, Tiemeier H, Hudziak JJ, et al. Television viewing and externalizing problems in preschool children: the Generation R Study. *Arch Pediatr Adolesc Med.* 2012;166(10):919–925. PubMed doi:10.1001/archpediatrics.2012.653
91. Hamer M, Stamatakis E, Mishra G. Psychological distress, television viewing, and physical activity in children aged 4 to 12 years. *Pediatrics.* 2009;123(5):1263–1268. PubMed doi:10.1542/peds.2008-1523
92. Tin SP, Ho DS, Mak KH, Wan KL, Lam TH. Association between television viewing and self-esteem in children. *J Dev Behav Pediatr.* 2012;33(6):479–485. PubMed doi:10.1097/DBP.0b013e31825ab67d
93. Robertson LA, McAnally HM, Hancox RJ. Childhood and adolescent television viewing and antisocial behavior in early adulthood. *Pediatrics.* 2013;131(3):439–446. PubMed doi:10.1542/peds.2012-1582
94. Mitrofan O, Paul M, Spencer N. Is aggression in children with behavioural and emotional difficulties associated with television viewing and video game playing? A systematic review. *Child Care Health Dev.* 2009;35(1):5–15. PubMed doi:10.1111/j.1365-2214.2008.00912.x

95. Nathanson AI, Sharp ML, Aladé F, Rasmussen EE, Christy K. The relation between television exposure and theory of mind among preschoolers. *J Commun.* 2013;63(6):1088–1108. doi:10.1111/jcom.12062
96. Duch H, Fisher EM, Ensari I, et al. Association of screen time use and language development in Hispanic toddlers: a cross-sectional and longitudinal study. *Clin Pediatr.* 2013;52(9):857–865. PubMed doi:10.1177/0009922813492881
97. van Egmond-Fröhlich AW, Weghuber D, de Zwaan M. Association of symptoms of attention-deficit/hyperactivity disorder with physical activity, media time, and food intake in children and adolescents. *PLoS One.* 2012;7(11):e49781. PubMed doi:10.1371/journal.pone.0049781
98. Bener A, Al-Mahdi HS, Ali AI, Al-Nufal M, Vachhani PJ, Tewfik I. Obesity and low vision as a result of excessive Internet use and television viewing. *Int J Food Sci Nutr.* 2011;62(1):60–62. doi:10.3109/09 637486.2010.495711
99. Cespedes EM, Gillman MW, Kleinman K, Rifas-Shiman SL, Redline S, Taveras EM. Television viewing, bedroom television, and sleep duration from infancy to mid-childhood. *Pediatrics.* 2014;133(5):e1163–e1171. PubMed doi:10.1542/peds.2013-3998
100. Marinelli M, Sunyer J, Alvarez-Pedrerol M, et al. Hours of television viewing and sleep duration in children: a multicenter birth cohort study. *JAMA Pediatr.* 2014;168(5):458–464. PubMed doi:10.1001/jamapediatrics.2013.3861
101. Mo-Suwan L, Nontarak J, Aekplakorn W, Sathannoppakao W. Computer game use and television viewing increased risk for overw8 among low activity girls: Fourth Thai National Health Examination Survey 2008–2009. *Int J Pediatr.* 2014;2014:364702.
102. Shan Z, Deng G, Li J, Li Y, Zhang Y, Zhao Q. Correlational analysis of neck/shoulder pain and low back pain with the use of digital products, physical activity and psychological status among adolescents in Shanghai. *PLoS One.* 2013;8(10):e78109.
103. Briggs AM, Straker LM, Bear NL, Smith AJ. Neck/shoulder pain in adolescents is not related to the level or nature of self-reported physical activity or type of sedentary activity in an Australian pregnancy cohort. *BMC Musculoskelet Disord.* 2009;10:87.
104. Barnett LM, Hinkley T, Okely AD, Hesketh K, Salmon J. Use of electronic games by young children and fundamental movement skills? *Percept Mot Skills.* 2012;114(3):1023–1034.
105. Anderson CA, Shibuya A, Ihori N, et al. Violent video game effects on aggression, empathy, and prosocial behavior in eastern and western countries: a meta-analytic review. *Psychol Bull.* 2010;136(2):151–173.
106. Brunborg GS, Mentzoni RA, Froyland LR. Is video gaming, or video game addiction, associated with depression, academic achievement, heavy episodic drinking, or conduct problems? *J Behav Addict.* 2014;3(1):27–32.
107. Syvaola HJ, Tammelin TH, Ahonen T, Kankaanpää A, Kantomaa MT. The associations of objectively measured physical activity and sedentary time with cognitive functions in school-aged children. *PLoS One.* 2014;9(7):e103559.

108. Kuhn S, Lorenz R, Banaschewski T, et al. Positive association of video game playing with left frontal cortical thickness in adolescents. *PLoS One*. 2014;9(3):e91506.
109. Cheung AC, Slavin RE. How features of educational technology applications affect student reading outcomes: a meta-analysis. *Educational Research Review*. 2012;7(3):198–215.
110. Schneps MH, Thomson JM, Chen C, Sonnert G, Pomplun M. E-readers are more effective than paper for some with dyslexia. *PloS One*. 2013;8(9):e75634.
111. Burton CE, Anderson DH, Prater MA, Dyches TT. Video self-modeling on an iPad to teach functional math skills to adolescents with autism and intellectual disability. *Focus Autism Other Dev Disabl*. 2013;28:67–77.
112. Hamilton JR. The electronic lonely crowd- patterns and effects of electronic media usage among contemporary adolescents. *International Journal of Arts & Sciences*. 2012;5(3):165–174.
113. Mathers M, Canterford L, Olds T, Hesketh K, Ridley K, Wake M. Electronic media use and adolescent health and well-being: crosssectional community study. *Acad Pediatr*. 2009;9(5):307–314.
114. Levine SC, Ratliff KR, Huttenlocher J, Cannon J. Early puzzle play: a predictor of preschoolers' spatial transformation skill. *Dev Psychol*. 2012;48(2):530–542.
115. Taylor BM, Frye BJ, Maruyama GM. Time spent reading and reading growth. *Am Educ Res J*. 1990;27(2):351–362.
116. Tang YY, Posner MI, Rothbart MK. Meditation improves self-regulation over the life span. *Ann N Y Acad Sci*. 2014;1307:104–111.
117. Yoo YG, Lee IS. The effects of school-based Maum meditation program on the self-esteem and school adjustment in primary school students. *Glob J Health Sci*. 2013;5(4):14–27.
118. Mitchell JA, Bottai M, Park Y, Marshall SJ, Moore SC, Matthews CE. A prospective study of sedentary behavior and changes in the BMI distribution. *Med Sci Sports Exerc*. 2014;46:2244–52.
119. Tanaka C, Reilly JJ, Huang WY. Longitudinal changes in objectively measured sedentary behaviour and their relationship with adiposity in children and adolescents: systematic review and evidence appraisal. *Obes Rev*. 2014;15:791–803.
120. Dowd KP, Harrington DM, Hannigan A, Donnelly AE. Light intensity physical activity is associated with adiposity in adolescent females. *Med Sci Sports Exerc*. 2014;46:295–300.
121. Hardy LL, Dobbins TA, Denney-Wilson EA, Okely AD, Booth ML. Sedentariness, small-screen recreation, and fitness in youth. *Am J Prev Med*. 2009;36(2):120–125.
122. Holmstrup M, Fairchild T, Keslacy S, Weinstock R, Kanaley J. Multiple short bouts of exercise over 12-h period reduce glucose excursions more than an energy-matched single bout of exercise. *Metabolism*. 2014;63(4):510–519.
123. Carson V, Stone M, Faulkner G. Patterns of sedentary behavior and weight status among children. *Pediatr Exerc Sci*. 2014;26(1):95–102.
124. Lopes L, Santos R, Pereira B, Lopes VP. Associations between sedentary behavior and motor coordination in children. *Am J Hum Biol*. 2012;24(6):746–752.

125. Trevelyan FC, Legg SJ. The prevalence and characteristics of back pain among school children in New Zealand. *Ergonomics*. 2010;53(12):1455–1460.
126. Straker LM, O’Sullivan PB, Smith AJ, Perry MC. Relationships between prolonged neck/shoulder pain and sitting spinal posture in male and female adolescents. *Man Ther*. 2009;14(3):321–329.
127. Murphy S, Buckle P, Stubbs D. Classroom posture and selfreported back and neck pain in schoolchildren. *Applied ergonomics*. 2004;35(2):113–120.
128. Faulkner G, Carson V, Stone M. Objectively measured sedentary behaviour and self-esteem among children. *Mental Health and Physical Activity*. 2014;7(1):25–29.
129. Chau JY, Grunseit AC, Chey T, et al. Daily sitting time and all-cause mortality: a meta-analysis. *PLoS One*. 2013;8(11):e80000.
130. Aitchison J. The statistical analysis of compositional data. *Journal of the Royal Statistical Society. Series B (Methodological)*. 1982:139– 177.
131. van Grieken A, Ezendam NP, Paulis WD, van der Wouden JC, Raat H. Primary prevention of overweight in children and adolescents: a meta-analysis of the effectiveness of interventions aiming to decrease sedentary behaviour. *Int J Behav Nutr Phys Act*. 2012;9:61.